



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Foreign and insular statistical reports of countries and cities—Yearly and monthly.

AFRICA—*Lourenço Marques*.—Month of February, 1902. Estimated population, 7,000. Total number of deaths, 21, including 3 from tuberculosis.

FRANCE—*Rouen*.—Month of March, 1902. Estimated population, 116,316. Total number of deaths, 287, including diphtheria, 4; enteric fever, 1; measles, 2; whooping cough, 1; smallpox, 1, and 60 from tuberculosis.

GERMANY—*Hanover*.—Month of February, 1902. Estimated population, 241,849. Total number of deaths, 279, including 22 from infectious diseases.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 76 great towns in England and Wales during the week ended April 12, 1902, correspond to an annual rate of 17.4 per 1,000 of the aggregate population, which is estimated at 14,862,909.

London.—One thousand five hundred and twenty-three deaths were registered during the week, including measles, 64; scarlet fever, 10; diphtheria, 23; whooping cough, 41; enteric fever, 1; smallpox, 73, and diarrhea and dysentery, 12. The deaths from all causes correspond to an annual rate of 17.3 per 1,000. In Greater London 2,099 deaths were registered. In the "outer ring" the deaths included 3 from diphtheria, 5 from measles, 1 from scarlet fever, 18 from smallpox, and 4 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended April 12, 1902, in the 21 principal town districts of Ireland was 24.2 per 1,000 of the population, which is estimated at 1,092,322. The lowest rate was recorded in Queenstown, viz, 0.0, and the highest in Portadown, viz, 46.7 per 1,000. In Dublin and suburbs 206 deaths were registered, including diphtheria, 1; enteric fever, 2; measles, 3; whooping cough, 2, and 41 from tuberculosis.

Scotland.—The deaths registered in 8 principal towns during the week ended April 12, 1902, correspond to an annual rate of 22.1 per 1,000 of the population, which is estimated at 1,679,923. The lowest mortality was recorded in Leith, viz, 13.9, and the highest in Greenock, viz, 26.5 per 1,000. The aggregate number of deaths registered from all causes was 715, including diphtheria, 2; measles, 10; scarlet fever, 8, smallpox, 2, and 21 from whooping cough.

GIBRALTAR.—Two weeks ended April 6, 1902. Estimated population, 27,460. Total number of deaths, 27, including 7 from measles.

JAMAICA—*Port Antonio*.—Two weeks ended April 19, 1902. Estimated population not reported. One death reported. No contagious diseases.

JAPAN—*Nagasaki*.—Ten days ended March 31, 1902. Estimated population, 131,700. Total number of deaths not reported. One death from enteric fever reported.

MALTA.—Four weeks ended April 12, 1902. Estimated population, 183,679. Total number of deaths, 302, including diphtheria, 4, and 6 from enteric fever.

ST. HELENA.—Two weeks ended March 22, 1902. Estimated population, 3,342. Total number of deaths, 10. No contagious diseases.

SPAIN—*Cadiz*.—Month of March, 1902. Estimated population, 70,177. Total number of deaths, 254, including diphtheria, 1; enteric fever, 5; measles, 16, and 28 from tuberculosis.

Valencia.—Two weeks ended April 15, 1902. Estimated population, 204,000. Total number of deaths, 177, including enteric fever, 3, and 6 from smallpox.

SWITZERLAND.—Reports for the two weeks ended April 5, 1902, from 18 cities and towns having an aggregate estimated population of 770,000, show a total of 587 deaths, including diphtheria, 7; enteric fever, 2; measles, 9; scarlet fever, 1; whooping cough, 17, and 77 from phthisis pulmonalis.

WEST INDIES—*St Thomas*.—Six weeks ended April 11, 1902. Estimated population, 11,012. Total number of deaths, 43. No deaths from contagious diseases.